Research Note

Poster 3: The acoustic analysis of vowel productions pre- and post-lingual frenectomy

Zuleica Camargo

Luciana Oliveira

Maria Augusta Forte Svicerio

Irene Queiroz Marchesan (*CEFAC, Brasil*)

Sandra Madureira (*Pontifical Catholic University of Sao Paulo*)

Suggested Citation

Camargo, Z., et al. (2016). Poster 3: The acoustic analysis of vowel productions pre- and post-lingual frenectomy. *International Journal of Orofacial Myology*, 42(1), 55-60.

DOI: <https://doi.org/10.52010/ijom.2016.42.1.7>

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

The views expressed in this article are those of the authors and do not necessarily reflect the policies or positions of the International Association of Orofacial Myology (IAOM). Identification of specific products, programs, or equipment does not constitute or imply endorsement by the authors or the IAOM. The journal in which this article appears is hosted on [Digital Commons](https://digitalcommons.elsevier.com/), an Elsevier platform.

POSTER 3: THE ACOUSTIC ANALYSIS OF VOWEL PRODUCTIONS PRE- AND POST- LINGUAL FRENECTOMY

ZULEICA CAMARGO, PH.D., SLP, LUCIANA OLIVEIRA, MA., SLP,
MARIA AUGUSTA FORTE SVICERO, MA., SLP,
IRENE MARCHESAN, PH.D., SLP
SANDRA MADUREIRA, PH.D., SLP

Awarded: Certificate of Excellence at the 2016 IAOM Convention

ABSTRACT

Introduction: Subjects with altered lingual frenulum may have difficulty speaking, chewing, and swallowing. Acoustic analysis allows for inferring the mechanisms of speech production, such as the tongue position on the horizontal and vertical axes and the jaw position. **Aim:** The aim is to compare formant frequencies (F1 and F2) for seven oral vowels in Brazilian Portuguese (BP), when produced by individuals who had a lingual frenectomy, pre- and post-surgical periods (a minimum six-month interval). **Methods:** There were seven subjects ranging in age from 6y3m to 14y7m, who were assessed and diagnosed with lingual frenulum alteration and the need for frenectomy was indicated, were selected and participated in two collections of speech data: (a) pre-surgery and (b) post-surgery. The speech samples consisted of words containing the seven vowels of BP which were acoustically analyzed in the software PRAAT, and the F1 and F2 measures were taken. The data were statistically analyzed (ANOVA) for the different vowels in the moments (a) and (b). **Results:** Acoustic measurements revealed significant differences between the pre-and post-surgical period (six months) when compared to F1 for vowels [a] ($p = 0.018$), [e] ($p = 0.018$) and [u] ($p = 0.049$) and F2 for vowels [e] ($p = 0.026$) and [u] ($p < 0.001$). **Conclusion:** F1 and F2 values indicated that the movement of the tongue improved after lingual frenectomy, revealing greater anterior mobilization but limited height range.

KEYWORDS: lingual frenulum, speech disorders, vowel, acoustic analysis

INTRODUCTION

Ankyloglossia or the deviation of the lingual frenulum has been reported to cause changes in tongue movement, thus impairing food functions and affecting speech (Chu & Blom, 2009; Obladen, 2009). Despite being a common condition, it is still not well defined in terms of diagnosis, treatment and prognosis.

With regards to frenulum alteration, we can see different types of alteration, for example, the lingual frenulum can be attached in the anterior aspect of the tongue near the tongue tip, and/or the length of the frenulum can be short. The anterior frenulum is known to affect the production of speech sounds, especially those that require refined movement of the tongue tip and body.

When the frenulum is short, it has a big impact upon the posture of the tongue, which remains on the floor of the mouth, and less impact on the movements of the tongue. Therefore, in such a condition, speech is less affected. In such cases, it is common that speech is produced utilizing a narrower range of motion by the articulators. When the frenulum is short and attached near the tip of the tongue, it generates a greater impact on speech, since the whole tongue remains on the mouth floor, with limited movements by the anterior part of the tongue (Marchesan, 2004).

Figure 1. Photographic records of pre-frenectomy (a) and post-frenectomy (b) in 4 cases of lingual frenectomy

In our clinical practice, we have observed that, the most common deviations found in produced by patients with an abnormal lingual frenulum, is the alveolar tap [r] (Camargo et al., 2013) and the alveolar fricatives [s] [z] productions. Other characteristics that affect articulation due to the limited tongue movement have to do with compensatory strategies or adaptations of jaw movements (e.g. excessive lateralization and anteriorization), and increased salivation, which can cause inaccuracies in speech production.

Although ankyloglossia is still subject to much controversy, surgical interventions have been proposed since ancient Greek medicine. One possible type of corrective surgery is lingual frenectomy, which is considered to be a safe and effective procedure for releasing the tongue-tie. This revision can lead to improved lingual function (Acevedo et al., 2010; Knox, 2010). The most common indicators of ankyloglossia are speech/articulation problems (64%), restricted movement (18%) and lactation problems/nutrition (8%) (Hall & Renfrew, 2005).

As for the results of surgery, improvement in mobility of the tongue and in the articulation of speech is the most reported effect (Klockars & Pitkaranta 2009; Obladen, 2009; Tuli & Singh, 2010). In Brazil, speech problems are very common in later ages, because the lingual frenectomy is not performed in early childhood. This can cause alteration and compensation of function related to the oral sensory-motor system, especially those that affect speech production. In this study, from the pre-operative period to the follow-up, we found that with the surgical release of the frenulum, the subjects' tongue movement amplitude increased, which yielded improved speech articulation, as well as an increase in the functional efficiency of the oral musculature for feeding functions, such as chewing and swallowing improved. During the immediate postoperative period, we observed improved speech production, for some adult patients. In some cases, however, change in speech production was not noticed in the early post frenectomy stage by random listeners or even the therapist, even though the extent of the functional articulation was observed to be significantly better.

Detailed characterization of the correlation between acoustic and perceptual characteristics of speech in subjects with abnormal lingual frenulum can assist in the process of treatment planning, therapeutic procedures and the evaluation process. There are no validated assessment protocols for treating patients with a disordered frenulum that have been used by medical professionals. In essence, there are no set ways to evaluate and interpret the results proposed. Marchesan (2004; 2005; 2010) has described speech problems as resulting from the alterations of the lingual frenulum and proposed protocols for evaluating this structure. The acoustic analysis, especially the characterization of the formant frequency of vowels, allows one to detail the mechanisms of the speech production, by inferring, from the acoustic data, the positioning of the tongue in the horizontal and vertical axes. Concerning this aspect, the aim in this study was to compare, from acoustic point of view, the formant frequencies (F1 and F2) for the seven oral vowels of Brazilian Portuguese Language (BP), when produced by the subject both pre- and post-

lingual frenotomy (a minimum six-month interval).

MATERIAL AND METHODS

Seven subjects, 06y 3m - 14y 07m, assessed and diagnosed with lingual frenulum alteration, without language and/or hearing disorders, were selected from patients attended by a Non-Government Organization. They participated in two sessions of speech data collection (audio recording).

The recordings were done in an acoustic environment, with the help of the BCA 2000 mixer (Behringer), Sound Forge software and AKG C140 headset microphone at two distinct moments: (a) pre- surgery and (b) post-surgery (Figure 1). The criteria to be included were: (a) diagnosed lingual frenulum alteration, frenectomy indicated (pre-surgical planning), and (b) attended over six months of speech therapy post-surgical period. The subjects were clinically assessed by the lingual frenulum protocol, which evaluates lingual

Table1. F1 and F2 frequency values (mean and standard deviation-SD) for the BP oral vowels produced by speakers before and after frenectomy

VOWEL	STAGE	F1			F2		
		Mean (HZ)	SD	p value	Mean (HZ)	SD	p value
[a]	Pre frenectomy	1020.71	162.73	0.018*	2330.52	724.35	0.456
	Post-frenectomy	1091.48	201.32		2340.71	912.85	
[ɛ]	Pre frenectomy	653.14	165.54	0.449	2705.76	261.82	0.026*
	Post-frenectomy	658.19	124.75		2837.10	265.82	
[e]	Pre frenectomy	467.83	69.97	0.018*	2972.67	436.35	0.363
	Post-frenectomy	423.52	99.57		3024.48	397.11	
[i]	Pre frenectomy	375.24	83.65	0.111	3166.48	389.01	0.063
	Post-frenectomy	349.57	68.72		3036.48	276.74	
[ɔ]	Pre frenectomy	653.24	141.00	0,059	1143.86	178.10	0.126
	Post-frenectomy	615.24	78.95		1094.43	242.94	
[o]	Pre frenectomy	482.45	72.21	0.101	891.83	135.83	0.101
	Post-frenectomy	472.33	65.01		936.29	110.39	
[u]	Pre frenectomy	421.62	95.00	0.049*	700.14	128.39	<0.001*
	Post-frenectomy	385.48	64.59		901.76	178.00	

aspects (Marchesan, 2010) and those of the possibilities of movement, as well as their possible interference in the functions of speech, breathing, swallowing and chewing.

The speech samples consisted of words inserted within a carrier sentence, containing the seven oral vowels of the BP. The vowel sounds were acoustically analyzed in PRAAT, a free access software (available at <http://www.fon.hum.uva.nl/praat/>). Formant frequencies (F1 and F2) values were obtained in Hz. The data were submitted to the CartaFormantesLogDoc.praat script, developed by Eduardo Velázquez (adaptation from Mietta & Lennes, 2004), in order to generate the vowel charts for both pre- and post- surgical data. Data were also compared to the International Phonetic Alphabet (IPA) description of Brazilian Portuguese (Barbosa & Albano, 2004).

Statistical Analysis (ANOVA) was also conducted to compare average frequencies of F1 and F2 for the different vowels at moments (a) and (b), with significance value established at $p < 0.05$. The research project was approved by the CEFAC Ethics Committee: 082/11.

RESULTS

The F1 and F2 formant frequency values and their standard deviation values for pre- and post-surgical lingual frenectomy are displayed in Table 1. The vowel charts, and their distribution of F1 and F2 values are presented in Figures 2 and 3 for pre- and post- surgical stages, respectively.

DISCUSSION

The findings support the assumption that lingual frenectomy increases the mobility of the tongue, making possible the restoration of oral functions previously described (Klockars, Pitkaranta, 2009; Aras et al., 2010; Acevedo et al., 2010; Knox, 2010; Tuli & Singh, 2010). However, the motor and timing controls of speech movements are not naturally restored after surgery and this condition mainly affects the adequacy of the speech production vowel patterns (Gregio et al., 2006; Ball & Gibbon, 2002). Access to acoustic analysis helps to identify different patterns of sound production in the pre- and post-surgical stages, so that the sounds classically described as "inaccurate" in the clinical evaluation, especially the vowels, can be analyzed to reveal the peculiarities in their acoustic structure.

Figure 2. Vowel chart (F1 and F2 distributions) for the BP oral vowels produced by speakers before frenectomy

Figure 3. Vowel chart (F1 and F2 distributions) for the BP oral vowels produced by speakers after frenectomy

Figure 4. Vowel chart (F1 and F2 distributions) for the BP oral vowels produced by children without lingual frenulum disorders (source: acoustic measures from Oliveira and Camargo (2016) p. 110)

The observed changes in vowel formant frequencies values were interpreted as revealing a tendency towards greater anteriorization of the tongue. This included the production of vowels, including the high back vowels, over a six-month period after the lingual frenectomy. However, the limited excursion of the tongue in the anterior-posterior axis persisted, as we can observe in Figures 2 and 3.

The vowel charts (figures 2 and 3) enabled us to observe the overlapping of vowel spaces, pre – frenotomy stage (a) and post-frenotomy – stage (b). At stage (b) partial excursion of the anterior vowels [i], [e] and [ε] spaces, but they are still overlapped, suggesting that the excursion in height axis was not totally matched. The same can be indicated for posterior vowels, despite the fact that the vowel spaces are a little more differentiated post-operatively, there is still some evidence of limited excursion of the tongue body in the posterior vowels.

In relation to F1 values, the most significant changes were found to occur in [a], [e] and [u]. For [a], when compared to Oliveira & Camargo (2016), the data in our study indicated that the tongue body (and/or the jaw) was still lower. It is worth mentioning that we also found more tongue raising in the post-operative stage, for the vowels [e] and, especially, for [u]. For F2 mean and

dispersion values, as displayed in Table 1, the most significant changes were found to occur in the group of anterior vowels for [ε], which had a tendency to occur with the tongue in a more anterior tongue posture. For back vowel [u], changes in F2 values also found a tendency towards a more anterior tongue posture. Although statistically significant differences for F2 average values were not found for vowel [a], dispersion values indicate a tendency towards an anterior tongue posture. When comparing F1 and F2 values found in this study to the ones mentioned in studies of BP vowels produced by a population without speech alterations- Figure 4 (Oliveira & Camargo 2016), it is notable that some of the values measured in this study do not fit the expectations for age and gender.

The results in our study were interpreted as showing a tendency, over a minimum six-month interval after lingual frenectomy, for greater anterior-wise mobilization of the tongue in the production of vowels, which includes both the back and the high vowels. Furthermore, it also revealed continued limited extension of the tongue in the anterior-posterior axis, especially for back vowels.

Based on that, speech therapy becomes a relevant and complementary treatment procedure. Because management of lingual frenulum alteration is still controversial among professionals (Obladen, 2009), the inclusion of protocols for evaluation and management of various alterations of the lingual frenulum may be beneficial. In reality, this outcome becomes more evident when considering that surgery is not commonly performed in young kids. In this case, acoustic analysis could prove to be an important tool to follow up the speech adaptations developed after frenectomy.

CONCLUSIONS

Changes in formant frequencies (F1 and F2) indicated that the range of motion and the function of the tongue improved over a period of six months after lingual frenectomy, revealing greater anterior mobilization, given speech therapy, although some limited height range was found to persist. The data reinforced the complexity of tongue (and jaw) movements during speech and the effects of frenectomy on speech production. Guidelines to speech rehabilitation can also be drawn from the discussed data.

CONTACT AUTHOR:

Zuleica Camargo, Ph.D., SLP

Integrated Acoustic Analysis and Cognition
Laboratory, Pontifical Catholic University of São
Paulo PUC-SP, Rua Monte Alegre, 969- Rooms
505-507 – 5th floor CEP: 05014-001 - São Paulo -
SP- Brazil
Business telephone number: +55.11.36708333
Fax number: +55.11.36708374
e-mail: zcamargo@pucsp.br

Luciana Oliveira, MA., SLP

CEFAC Heath & Education, São Paulo, Brazil.

Maria Augusta Forte Svicero, MA., SLP

CEFAC Heath & Education, São Paulo, Brazil.

Irene Marchesan, Ph.D., SLP

CEFAC Heath & Education, São Paulo, Brazil.

Sandra Madureira, Ph.D., SLP

Pontifical Catholic University of São Paulo PUC-SP
São Paulo, Brazil.

REFERENCES

- Aras, M. H., Göregen, M., Güngörmüş, M. & Akgül, H. M. (2010). Comparison of Diode Laser and Er: YAG Lasers in the Treatment of Ankyloglossia. *Photomedicine and laser surgery*, 28(2), 173-177.
- Acevedo, A. C., da Fonseca, J. A. C., Grinham, J., Doudney, K., Gomes, R. R., de Paula, L. M. & Stanier, P. (2010). Autosomal-dominant ankyloglossia and tooth number anomalies. *Journal of dental research*, 89(2), 128-132.
- Ball, M. J. & Gibbon, F. E. (2002). *Vowel disorders*. Butterworth-Heinemann.
- Barbosa, P. A. & Albano, E. C. (2004). Brazilian Portuguese. *Journal of the International Phonetic Association*, 34(02), 227-232.
- Camargo, Z. A., Marchesan, I. Q., Oliveira, L. R., Svicero, M. A. F., Pereira, L. C. K. & Madureira, S. (2013). Lingual frenectomy and alveolar tap production: An acoustic and perceptual study. *Logopedics Phoniatrics Vocology*, 38(4), 157-166.
- Chu, M. W. & Bloom, D. C. (2009). Posterior ankyloglossia: A case report. *International journal of pediatric otorhinolaryngology*, 73(6), 881-883.
- Gregio, F. N., Gama-Rossi, A., Madureira, S. & Camargo, Z. (2006). Modelos teóricos de produção e percepção da fala como um sistema dinâmico. *Rev. CEFAC*, 8(2), 244-247.
- Hall, D. M. B. & Renfrew, M. J. (2005). Tongue tie. *Archives of disease in childhood*, 90(12), 1211-1215.
- Klockars, T. & Pitkäranta, A. (2009). Pediatric tongue-tie division: indications, techniques and patient satisfaction. *International journal of pediatric otorhinolaryngology*, 73(10), 1399-1401.
- Knox, I. (2010). Tongue tie and frenotomy in the breastfeeding newborn. *NeoReviews*, 11(9), e513-e519.
- Lennes, M. (2004). Praat-opas. Internet-versio luettavissa osoitteessa: <http://www.helsinki.fi/puhetieteet/atk/praat/praat.pdf>.
- Marchesan, I.Q. (2004). Lingual frenulum: Classification and speech interference. *The International journal of orofacial myology: official publication of the International Association of Orofacial Myology*, 30, 31-38.
- Marchesan, I.Q. (2005). Lingual Frenulum: Quantitative evaluation proposal. *International Journal of Orofacial Myology*, 31, 39-48.
- Marchesan, I.Q. (2010). Protocolo de avaliação do frênulo da língua. *Rev Cefac*, 12(6), 977-99.
- Obladen, M. (2009). Much ado about nothing: Two millenia of controversy on tongue-tie. *Neonatology*, 97(2), 83-89.
- Oliveira, L.R. & Camargo, Z. (2016). Análise acústica comparativa das vogais orais entre crianças respiradoras orais e nasais. In: Camargo, Z. Fonética clínica: 20 anos de LIAAC. São José dos Campos: Pulso. p105-19.
- Tuli, A. & Singh, A. (2010). Monopolar diathermy used for correction of ankyloglossia. *Journal of Indian Society of Pedodontics and Preventive Dentistry*, 28(2), 130