Research Article

Case presentation: Dental treatment with PAP for ALS patient

Takeshi Kikutani (*The Rehabilitation Clinic for Speech and Swallowing Disorders, the Nippon Dental University Tokyo Hospital*)

Fumiyo Tamura

Keiko Nishiwaki

Suggested Citation

Kikutani, T., et al. (2006). Case presentation: Dental treatment with PAP for ALS patient. *International Journal of Orofacial Myology*, 32(1), 32-35.

DOI: <https://doi.org/10.52010/ijom.2006.32.1.3>

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

The views expressed in this article are those of the authors and do not necessarily reflect the policies or positions of the International Association of Orofacial Myology (IAOM). Identification of specific products, programs, or equipment does not constitute or imply endorsement by the authors or the IAOM. The journal in which this article appears is hosted on [Digital Commons](https://digitalcommons.elsevier.com/), an Elsevier platform.

CASE PRESENTATION: DENTAL TREATMENT WITH PAP FOR ALS PATIENT

Takeshi Kikutani, D.D.S., Ph.D., Fumiyo Tamura, D.D.S., Ph.D.,
Keiko Nishiwaki, S.L.P.

ABSTRACT

The aim of this study was to evaluate the effect of Palatal Augmentation Prosthesis (PAP) for an Amyotrophic Lateral Sclerosis (ALS) patient. The patient's palatogram was taken during swallowing to assess her lingual function. A PAP was provided to assist her with feeding and swallowing functions. The patient's lingual pressure strength showed increases, and she was able to feed well using PAP. This device was useful for assisting lingual dysfunctions due to various diseases.

Key Words: Orofacial Myofunctional Disorders, Amyotrophic Lateral Sclerosis, Palatal Augmentation Prosthesis, Lingual Function

INTRODUCTION

Palatal Augmentation Prosthesis (PAP) is a device used for oral cancer patients (Wheeler, Logeman, Rosen, 1980; Logemann, Kahrias, Hurstm Davis, Krugler 1989) and stroke patients (Ono T, Hamamura, Honda, Nokubi, 2005) with tongue dysfunctions. The aim of this study is to evaluate the effect of PAP for an Amyotrophic Lateral Sclerosis patient.

The tongue plays an important role in oral functions. The tongue moves toward the palate while changing its shape during mastication or swallowing. This function will be disturbed when the tongue can not reach the palate due to atrophy or glossectomy. Consequently, PAP works to support the tongue in touching the palate and recovering lingual function by placing a plate on the palate.

Mechanism of PAP

An increased oral cavity due to alteration of relative oral capacity caused by atrophy or removal of the tongue disturbs the contact of tongue to the palate during the preparatory, oral and pharyngeal stages. Thus, dental treatment with PAP is a prosthetic approach for supporting feeding/swallowing functions. Inadequate tongue contact with the palate may provoke backward movement of tongue base due to declined lingual function, which tends to prevent the tongue movement

toward the palate. The purpose of PAP is also to reinforce these functions. This is an aspect of a functional approach for improving swallowing.

However, the previous studies reported not only PAP for tongue glossectomy cases but also for feeding functions of ALS patients. In addition, there was a paper about PAP for dysarthria (Esposito 2000). Thus, we have been trying to use PAP for ALS patients and evaluate the effectiveness.

Clinical Case: Amyotrophic Lateral Sclerosis patient

ALS, first described in 1869, is a progressive neurodegenerative disease that attacks nerve cells in the brain and spinal cord resulting in muscle weakness and atrophy. [There were reports which showed severe situation according to their life] (Sejvar, Holman, Bresee, Kochanek, 2005; Van den Berg JP, Katmilin, Linderman, Veldink, deVisser, Van der Graff, Van de Berg, LH, 2005). At the onset of ALS, the symptoms may be quite slight and overlooked frequently. The appearance of symptoms and the progression of the illness, the course of the disease are seen as follows: 1) muscle weakness in one or more of the following: hands, arms, legs or the muscles of breathing, swallowing or speech, and fasciculation, 2) cramping of muscles in the

hands and feet, 3) disability of the arms and legs, 4) difficulty in projecting the voice, 5) shortness of breath, difficulty in breathing and swallowing coordination.

Case Profile

This case is a 48-year-old female. Her major diagnosis was ALS with severe dysphagia dysfunction. The chief complaint was difficulty with the transport of the food bolus to the pharynx and inability to swallow. Oral conditions were dentate and tongue atrophy.

Palatogram on the PAP

A palatal plate as the base was fabricated (Figure 1). The patient's palatogram was taken during a dry swallow. Then we could observe whether or not the tongue touched the palate.

Figure 1. Taking a palatogram

At first, the patient touched her tongue toward the central part of plate during swallowing inadequately. Therefore we continued to reshape the base plate with dental wax until her tongue contacted the palate in a stable manner.

Fitting the PAP

The PAP was set in the palatal area (Figure 2). The thickness of the palatal plate of the PAP was increased to secure adequate contact of the tongue. Initially the patient complained that the apparatus was quite uncomfortable. However she adjusted to it rapidly with every day use in about one month. After this adjustment period her eating speed increased tremendously.

Figure 2. Fit the PAP

ASSESSMENT OF FOOD TEST - WITHOUT THE PAP

The effect of PAP was assessed by the Food Test. The Food Test was described by Ishida and Mukai (2002) as: first, the patient chews 4-grams of test food similar to oatmeal; next, the patient swallows it once; and then the patient opens the mouth to evaluate the amount of food residue on the tongue.

Figure 3. Food residue without PAP

In this case without PAP, there was food residue on the patient's tongue surface (Figure 3). This suggested that she could not push her tongue toward the palate adequately to clear the oral cavity.

ASSESSMENT OF FOOD TEST - WITH THE PAP

The efficacy of this apparatus is shown in Figure 4. Food residue decreased tremendously after she received the PAP.

Figure 4. No food residue with PAP

Effect of the PAP -Maximum lingual pressure

Lingual pressure pushing toward the palate was measured using a pressure sensor. The Handy manometer (ALNIC Co., JAPAN) (Hayashi, Tsuga, Hosokawa, Yoshida, Sato, Akagawa, 2002) was used for measurements of maximum lingual pressure. The lingual pressure against the bubble shaped sensors in contact with the hard palate was evaluated. Measurement was taken five times. The mean values and standard deviations were obtained by averaging the values for five measurements.

Contact author:

Takeshi Kikutani DDS, Ph.D. Associate Professor,
The Rehabilitation Clinic for Speech and Swallowing Disorders, the Nippon Dental University Tokyo Hospital
2-3-16 Fujimi Chiyoda-ku, Tokyo, Japan 102-8158
TEL; 81-3-3261-5511
FAX;81-3-3261-3924 Email<kikutani@tokyo.ndu.ac.jp>

About the Authors:

1. Takeshi Kikutani DDS, Ph.D. Associate Professor, The Rehabilitation Clinic for Speech and Swallowing Disorders, the Nippon Dental University Tokyo Hospital, Tokyo, Japan
2. Fumiyo Tamura DDS, Ph.D. Assistant Professor, The Rehabilitation Clinic for Speech and Swallowing Disorders, The Nippon Dental University Tokyo Hospital, Tokyo Japan
3. Keiko Nishiwaki S.L.P. The Rehabilitation Clinic for Speech and Swallowing Disorders, The Nippon Dental University Tokyo Hospital, Tokyo Japan

Figure 5. Effect of PAP for lingual pressure

According to this result, the lingual pressure with the PAP showed an increase compared to that without the PAP (Figure 5). It is proposed that this was the result of an improvement in lingual function because the tongue could work better during swallowing. In fact, the patient realized that her feeding function improved as there was less food residue after swallowing with this apparatus and she could also eat faster using this device.

CONCLUSION

PAP was useful and effective for an ALS patient who experienced deterioration of lingual function. The authors believe that dentists should contribute their knowledge and techniques to patients experiencing ALS.

REFERENCES

- Esposito SJ, Mitumono H, Shanks M.** (2000). Use of palatal lift and palatal augmentation prostheses to improve dysarthria in patients with amyotrophic lateral sclerosis: A case series. Journal Prostodontic Dentistry. 83: 90-8
- Hayashi R, Tsuga K, Hosokawa R, Yoshida M, Sato Y, Akagawa Y.** (2002). A novel handy probe for tongue pressure measurement, International Journal Prosthodontics. 15 : 385-388.
- Ishida R, Mukai Y.** (2002). New assessment method of swallowing function; Staged food test. Journal of Clinical Rehabilitation. 11(9): 820-824
- Logemann JA, Kahrilas PJ, Hurst P, Davis J, Krugler C.** (1989). Effects of intraoral prosthetics on swallowing in patients with oral cancer. Dysphagia. 4(2):118-20.
- Ono T, Hamamura M, Honda K, Nokubi T.** (2005) Collaboration of a dentist and speech-language pathologist in the rehabilitation of a stroke patient with dysarthria: a case study. Gerodontology. 22(2):116-9.
- Sejvar JJ, Holman RC, Bresee JS, Kochanek KD, Schonberger LB.** (2005) Amyotrophic lateral sclerosis mortality in the United States, 1979-2001. Neuroepidemiology. 25(3): 144-52.
- Van den Berg JP, Kalmijin S, Linderman E, Veldink JH, de Visser M, Van der Graaff MM, Wokke JH, Van de Berg LH** (2005). Multidisciplinary ALS care improves quality of life in patients with ALS. Neurology. 65(8): 1264-7.
- Wheeler RL, Logemann JA, Rosen MS.** (1980). Maxillary reshaping prostheses: effectiveness in improving speech and swallowing of postsurgical oral cancer patients. Journal Prosthetic Dentistry. 43(3):313-9.